

Foot-shock stimulation decreases the inhibitory action of atp on contractility and end-plate current of frog sartorius muscle

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Ayrat Usmanovich Ziganshin et al. Background and Objective: Inhibitory action of ATP on contractions of frog isolated sartorius muscle was shown earlier, while hydrocortisol prevented that effect of ATP. The aim of this study was to investigate whether ATP also inhibits the contractility of sartorius muscle on stress-induced animals. Materials and Methods: The influence of ATP on contractions and end-plate current (EPC) of sartorius muscles isolated from frogs exposed to electric foot-shock stimulation (EFSS) was analyzed. Results: In the muscles of frogs exposed to EFSS, the inhibitory effect of ATP on contractions was significantly decreased and on EPC was abolished. The level of cortisol was much higher in the blood obtained from EFSS animals comparing with control ones. Conclusion: The EFSS has inhibitory action on the effect of ATP on the frog sartorius muscle and that could be due to increase of cortisol blood level.

<http://dx.doi.org/10.3923/ijp.2018.1198.1202>

Keywords

ATP, Contractions, Cortisol, End-plate current, Foot-shock, Frog sartorius muscle

References

- [1] Grishin, S.N. and A.U. Ziganshin, 2013. Modulatory role of purines in neuromuscular transmission. Biochem. (Moscow) Suppl. Ser. A: Membr. Cell Biol., 7: 183-191.
- [2] Grishin, S.N., A.Y. Teplov, A.V. Galkin, A.M. Devyataev and A.L. Zefirov et al., 2006. Different effects of ATP on the contractile activity of mice diaphragmatic and skeletal muscles. Neurochem. Int., 49: 756-763.
- [3] Giniatullin, A.R., S.N. Grishin and R.A. Giniatullin, 2000. Effect of hydrocortisone on purine actions in the nerve-muscle preparation. Rossiiskii Fiziologicheskii Zhurnal Imeni IM Sechenova, 86: 1293-1299.
- [4] Passaquin, A.C., P. Lhote and U.T. Ruegg, 1998. Calcium influx inhibition by steroids and analogs in C2C12 skeletal muscle cells. Br. J. Pharmacol., 124: 1751-1759.
- [5] Chaban, V.V., E.A. Mayer, H.S. Ennes and P.E. Micevych, 2003. Estradiol inhibits ATP-induced intracellular calcium concentration increase in dorsal root ganglia neurons. Neuroscience, 118: 941-948.
- [6] Ziganshin, A.U., R.R. Kamaliev, S.N. Grishin, L.E. Ziganshina, A.L. Zefirov and G. Burnstock, 2005. The influence of hypothermia on P2 receptor-mediated responses of frog skeletal muscle. Eur. J. Pharmacol., 509: 187-193.
- [7] Ziganshin, A.U., R.R. Kamaliev, S.N. Grishin, B.A. Ziganshin and G. Burnstock, 2009. Interaction of hydrocortisone with ATP and adenosine on nerve-mediated contractions of frog skeletal muscle. Eur. J. Pharmacol., 607: 54-59.
- [8] Sokolova, E., S. Grishin, A. Shakirzyanova, M. Talantova and R. Giniatullin, 2003. Distinct receptors and different transduction mechanisms for ATP and adenosine at the frog motor nerve endings. Eur. J. Neurosci., 18: 1254-1264.

- [9] Grishin, S., A. Shakirzyanova, A. Giniatullin, R. Afzalov and R. Giniatullin, 2005. Mechanisms of ATP action on motor nerve terminals at the frog neuromuscular junction. *Eur. JNeurosci.*, 21: 1271-1279.
- [10] Han, J.Z., W. Lin and Y.Z. Chen, 2005. Inhibition of ATP-induced calcium influx in HT4 cells by glucocorticoidsInvolvement of protein kinase A. *Acta Pharmacol. Sin.*26: 199-204.
- [11] Rogers, K.M., C.A. Bonar, J.L. Estrella and S. Yang, 2002. Inhibitory effect of glucocorticoid on coronary artery endothelial function. *Am. J. Physiol.-Heart Circulat. Physiol.*, 283: H1922-H1928.
- [12] Di Virgilio, F., P. Chiozzi, S. Falzoni, D. Ferrari, J.M. Sanz, V. Venketaraman and O.R. Baricordi, 1998. Cytolytic P2X purinoceptors. *Cell Death Different.*, 5: 191-199.
- [13] Goulding, N.J., 2004. The molecular complexity of glucocorticoid actions in inflammation-a four-ring circus. *Curr. Opin. Pharmacol.*, 4: 629-636.